

Osteoporosis Enrollment Form

Patient Data	SSN #: XXX-XX- Address: Home Phone: Alternate Caregiver Name:	Birthdate: Cell Phone:	Known Allergies: City: Primary Language: Phone of Caregiver:		Zip:
Ins. Data		Group#:			Group#:
Diagnosis	□ M80.0 Osteoporosis with pathological fracture □ Other: □ M81.0 Age-related osteoporosis □ M81.8 Other Osteoporosis				
Clinical Information	To expedite prior authorization, please attach clinical office notes, DEXA Scan report, and any labs completed. Lowest Dexa T-score:Site:Date:/				
	Reason(s) for discontinuation: Medication Directions Ouantity Refills				
	Medication	Directions		Ouantity	Refills
	Medication ☐ Boniva 3mg/3ml PFS Kit	Infuse 3mg IV over 15-30 seconds every 3 m	nonths	Quantity #1 kit	Refills
	·				Refills
Medication	□ Boniva 3mg/3ml PFS Kit	Infuse 3mg IV over 15-30 seconds every 3 m Inject 2 syringes (210mg), one after the otl	her, subcutaneously	#1 kit	Refills
Medication	□ Boniva 3mg/3ml PFS Kit □ Evenity™ 105mg/1.17ml PFS □ Forteo 600mcg/2.4ml pen □ Mini Pen Needles for Forteo	Infuse 3mg IV over 15-30 seconds every 3 m Inject 2 syringes (210mg), one after the oth in separate areas once monthly Inject 20mcg subcutaneously one time daily	her, subcutaneously	#1 kit #2 syringes #1 pen	Refills
Medication	□ Boniva 3mg/3ml PFS Kit □ Evenity™ 105mg/1.17ml PFS □ Forteo 600mcg/2.4ml pen □ Mini Pen Needles for Forteo Injection	Infuse 3mg IV over 15-30 seconds every 3 m Inject 2 syringes (210mg), one after the oth in separate areas once monthly Inject 20mcg subcutaneously one time daily Use as directed	her, subcutaneously y	#1 kit #2 syringes #1 pen #100	Refills
	□ Boniva 3mg/3ml PFS Kit □ Evenity™ 105mg/1.17ml PFS □ Forteo 600mcg/2.4ml pen □ Mini Pen Needles for Forteo Injection □ Prolia 60mg/1ml PFS □ Reclast 5mg/100ml	Infuse 3mg IV over 15-30 seconds every 3 m Inject 2 syringes (210mg), one after the other in separate areas once monthly Inject 20mcg subcutaneously one time daily Use as directed Inject 60mg subcutaneously every 6 months	her, subcutaneously y s minutes, once	#1 kit #2 syringes #1 pen #100 #1 syringe	
	□ Boniva 3mg/3ml PFS Kit □ Evenity™ 105mg/1.17ml PFS □ Forteo 600mcg/2.4ml pen □ Mini Pen Needles for Forteo Injection □ Prolia 60mg/1ml PFS □ Reclast 5mg/100ml	Infuse 3mg IV over 15-30 seconds every 3 m Inject 2 syringes (210mg), one after the other in separate areas once monthly Inject 20mcg subcutaneously one time daily Use as directed Inject 60mg subcutaneously every 6 months Infuse 5mg (100ml) IV, over no less then 15 every year	her, subcutaneously y s minutes, once Patient Office Otl	#1 kit #2 syringes #1 pen #100 #1 syringe #1 vial her Need by Date:	
	□ Boniva 3mg/3ml PFS Kit □ Evenity™ 105mg/1.17ml PFS □ Forteo 600mcg/2.4ml pen □ Mini Pen Needles for Forteo Injection □ Prolia 60mg/1ml PFS □ Reclast 5mg/100ml Is patient new to this to Prescriber Name: Practice Name:	Infuse 3mg IV over 15-30 seconds every 3 m Inject 2 syringes (210mg), one after the other in separate areas once monthly Inject 20mcg subcutaneously one time daily Use as directed Inject 60mg subcutaneously every 6 months Infuse 5mg (100ml) IV, over no less then 15 every year	her, subcutaneously y s minutes, once Patient □ Office □ Otl DEA#: Contact:	#1 kit #2 syringes #1 pen #100 #1 syringe #1 vial her Need by Date:	d: □ Phone □ Fax □ Email
escriber Data	□ Boniva 3mg/3ml PFS Kit □ Evenity™ 105mg/1.17ml PFS □ Forteo 600mcg/2.4ml pen □ Mini Pen Needles for Forteo Injection □ Prolia 60mg/1ml PFS □ Reclast 5mg/100ml Is patient new to this t Prescriber Name: Practice Name: Address:	Infuse 3mg IV over 15-30 seconds every 3 m Inject 2 syringes (210mg), one after the other in separate areas once monthly Inject 20mcg subcutaneously one time daily Use as directed Inject 60mg subcutaneously every 6 months Infuse 5mg (100ml) IV, over no less then 15 every year	her, subcutaneously y s minutes, once Patient Office Otl DEA#: Contact: Phone:	#1 kit #2 syringes #1 pen #100 #1 syringe #1 vial her Need by Date:	d: □ Phone □ Fax □ Email Fax:
	□ Boniva 3mg/3ml PFS Kit □ Evenity™ 105mg/1.17ml PFS □ Forteo 600mcg/2.4ml pen □ Mini Pen Needles for Forteo Injection □ Prolia 60mg/1ml PFS □ Reclast 5mg/100ml Is patient new to this to Prescriber Name: Practice Name: Address: City: State	Infuse 3mg IV over 15-30 seconds every 3 m Inject 2 syringes (210mg), one after the other in separate areas once monthly Inject 20mcg subcutaneously one time daily Use as directed Inject 60mg subcutaneously every 6 months Infuse 5mg (100ml) IV, over no less then 15 every year	her, subcutaneously y s minutes, once Patient Office Otl DEA#: Contact: Phone: Email:	#1 kit #2 syringes #1 pen #100 #1 syringe #1 vial her Need by Date:	d: □ Phone □ Fax □ Email Fax:

fax referral to: 901-388-0407
RESTORE RX OSTEOPOROSIS ENROLLMENT FORM | MARCH 23, 2022 7:13 PM

phone: **877-388-050**7

www.restorerx.com